

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES


Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

**As rescanning documents *will not* correct images,
please do not report the images to the
Image Problem Mailbox.**



Chemical diversity in different extraction solvents.
Root extracts from *Daucus carota* (carrot), elicited
with 1 mM AgNO₃.
HPLC-profiles with UV detection at 254 nm.

Figure 11




Figure 1

*Soybean plants
diseased*




Figure 2




Figure 3






Figure 4




Effect of elicitation on the chemical composition of root exudates
of Brassica juncea.
HPLC-profiles with UV detection at 254 nm.

Figure 5




Effect of elicitation on the chemical composition of root exudates of *Datura metel*.
HPLC-profiles with UV detection at 254 nm.

Figure 6




Effect of elicitation on the chemical composition of root exudates
of Lupinus polyphyllus.
HPLC-profiles with UV detection at 254 nm.

Figure 7




Effect of elicitation on the chemical composition of root exudates
of *Melilotus medicaginoides*.
HPLC-profiles with UV detection at 254 nm.

Figure 8




Chemical diversity in different extraction solvents.
Root extracts from *Solanum melongena* (eggplant).
HPLC-profiles with UV detection at 254 nm.

Figure 9




Chemical diversity in different extraction solvents.
Root extracts from *Solanum melongena* (eggplant), elicited
with 1 mM Salicylic acid.
HPLC-profiles with UV detection at 254 nm.

Figure 10




Chemical diversity in different extraction solvents.
Root extracts from *Daucus carota* (carrot), elicited
with 1 mM AgNO₃.
HPLC-profiles with UV detection at 254 nm.

Figure 11




Effect of elicitation on chemical diversity of root extracts.
EtOAc phases of extracts from *Glycyne max* (soybean).
Total Ion Current of chromatograms scanned from 70 m/z to 400 m/z.
e - Elicitor peak

Figure 12




Effect of elicitation on chemical diversity of root extracts.
MeOH/MeCl₂ phases of extracts from *Daucus carota* (carrot).
HPLC-profiles with UV detection at 254 nm.

Figure 13




Effect of elicitation on chemical diversity of root extracts.
EtOAc phases of extracts from *Daucus carota* (carrot).
HPLC-profiles with UV detection at 254 nm.

Figure 14



Effect of elicitation on chemical diversity of root extracts.
EtOAc phases of extracts from *Lycopersicon esculentum* (tomato).
HPLC-profiles with UV detection at 254 nm.

Figure 15



Effect of UV irradiation on chemical diversity of root extracts.
EtOAc phases of extracts from *Lupinus polyphyllus* (lupine).
HPLC-profiles with UV detection at 254 nm.

Figure 16

1513

1515

1517

1519

1521

1523




Figure 17




Figure 18




Figure 19




Figure 20




Figure 21




Figure 22




Figure 23




Figure 24




Figure 25




Figure 26




Figure 27




Figure 28




Figure 29




Figure 30




Figure 31